size: 104x161 mm

## **ORAFIL-G**

## TEMPORARY FILLING MATERIAL



**Product Description**: Orafil-G is a paste consistency, radiopaque, temporary filling material which cures upon contact with oral fluids. It possesses high adhesive properties and therefore offers best possible closure of dental cavities

Composition: Zinc oxide, Calcium Sulphate, Accelerator, Flavouring Agents and Excipients.

Intended purpose/Intended use: It is intended to be used as a temporary filling material. Indications for use: Recommended for temporary restoration of cavities in the interval between tooth preparation and the permanent cementation of completed in lays.

Instructions for use: Use any suitable instrument to introduce the paste directly into wet cavity. Press and spread with the instrument to fill the cavity and adapt to the cavity walls. Remove any excess paste by wiping it off with a moist cotton pellet towards the margins of the cavity. Hardened paste can be removed with the use of rotary instrument or any suitable hand instrument. It is advised not to fill very deep cavities up to the base of cavity. The dispensed material shall not be placed back in container and the dose once applied and used shall not be reused.

Warnings: Avoid prolonged or repeated contact with skin (allergic contact dermatitis) and eyes. Avoid prolonged inhalation. Do not take internally. Contact with the skin may cause irritation, burns of hypersensitivity. If contact with skin occurs immediately wipe off thoroughly with cotton and alcohol and then wash well with soap and water. If skin rash and sensitization or other allergic reaction occurs discontinue use and seek medical attention. If the product comes into contact with the eyes, rinse thoroughly with fresh water and consult a physician. It should not be used with patients who have a history of severe allergic reaction to any of the components.

**Precautions:** In sensitive persons, the product may cause sensitization. If allergic reaction occurs remove the filling immediately and consult physician.

Contraindications: Do not apply directly on exposed pulp.

Storage Conditions: Store at temperature between 10°C(50°F) to 24°C(75°F). Close the cap carefully after use. Keep away from moisture. Shelf life not affected after opening if stored in a cool dry place.

Shelf Life: 3 years from the date of manufacturing.

## Presentation:

Ref: 11001

1x 40a Jar.

## Disposal Conditions:

To be disposed off based on the Prevailing rules of the country in which it is used.



stipulated area of applications. The user is responsible for testing the material for its suitability and use for any purpose not explicitly stated in this instruction sheet. Description and data constitute no warranty of attributes and are not binding.

For feedback, compliants and reporting of unexpected events contact customer care.